

28th International Mathematical Olympiad

Havana, Cuba

Day I

July 10, 1987

1. Let $p_n(k)$ be the number of permutations of the set $\{1, \dots, n\}$, $n \geq 1$, which have exactly k fixed points. Prove that

$$\sum_{k=0}^n k \cdot p_n(k) = n!.$$

(Remark: A permutation f of a set S is a one-to-one mapping of S onto itself. An element i in S is called a fixed point of the permutation f if $f(i) = i$.)

2. In an acute-angled triangle ABC the interior bisector of the angle A intersects BC at L and intersects the circum-circle of ABC again at N . From point L perpendiculars are drawn to AB and AC , the feet of these perpendiculars being K and M respectively. Prove that the quadrilateral $AKNM$ and the triangle ABC have equal areas.
3. Let x_1, x_2, \dots, x_n be real numbers satisfying

$$x_1^2 + x_2^2 + \dots + x_n^2 = 1.$$

Prove that for every integer $k \geq 2$ there are integers a_1, a_2, \dots, a_n , not all 0, such that $|a_i| \leq k - 1$ for all i and

$$|a_1x_1 + a_2x_2 + \dots + a_nx_n| \leq \frac{(k-1)\sqrt{n}}{k^n - 1}.$$

28th International Mathematical Olympiad

Havana, Cuba

Day II

July 11, 1987

4. Prove that there is no function f from the set of non-negative integers into itself such that $f(f(n)) = n + 1987$ for every n .
5. Let n be an integer greater than or equal to 3. Prove that there is a set of n points in the plane such that the distance between any two points is irrational and each set of three points determines a non-degenerate triangle with rational area.
6. Let n be an integer greater than or equal to 2. Prove that if $k^2 + k + n$ is prime for all integers k such that $0 \leq k \leq \sqrt{n/3}$, then $k^2 + k + n$ is prime for all integers k such that $0 \leq k \leq n - 2$.